



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Lexis and grammar of mitochondrial RNA processing in Trypanosomes

Citation for published version:

Aphasizheva, I, Alfonzo, J, Carnes, J, Cestari, I, Cruz-Reyes, J, Göringer, HU, Hajduk, S, Lukeš, J, Madison-Antenucci, S, Maslov, DA, McDermott, SM, Ochsenreiter, T, Read, LK, Salavati, R, Schnauffer, A, Schneider, A, Simpson, L, Stuart, K, Yurchenko, V, Zhou, ZH, Zíková, A, Zhang, L, Zimmer, S & Aphasizhev, R 2020, 'Lexis and grammar of mitochondrial RNA processing in Trypanosomes', *Trends in Parasitology*, vol. 36, no. 4, pp. 337-355. <https://doi.org/10.1016/j.pt.2020.01.006>

Digital Object Identifier (DOI):

[10.1016/j.pt.2020.01.006](https://doi.org/10.1016/j.pt.2020.01.006)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Trends in Parasitology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Trends in Parasitology

Lexis and Grammar of Mitochondrial RNA Processing in Trypanosomes

--Manuscript Draft--

Manuscript Number:	
Article Type:	Review
Keywords:	Trypanosoma; mitochondria; kinetoplast; RNA editing; RNA decay; polyadenylation; translation.
Corresponding Author:	Inna Afasizheva, Ph.D. Boston University Medical Center Boston, MA UNITED STATES
First Author:	Inna Afasizheva, Ph.D.
Order of Authors:	Inna Afasizheva, Ph.D. Juan Alfonso Jason Carnes Igor Cestari Jorge Cruz-Reyes H. Ulrich Göringer Stephen Hajduk Julius Lukeš Susan Madison-Antenucci Dmitri A. Maslov Suzanne M. McDermott Torsten Ochsenreiter, Laurie K. Read Reza Salavati Achim Schnauffer André Schneider Larry Simpson Kenneth Stuart Vyacheslav Yurchenko Z. Hong Zhou Alena Zíková Liye Zhang Sara Sara Zimmer Ruslan Aphasizhev
Abstract:	Parasitic protist <i>Trypanosoma brucei</i> sp. cause African human and animal trypanosomiasis. These hemoflagellates belong to the class Kinetoplastea, a taxon distinguished by the presence of a kinetoplast. This nucleoprotein body contains interlinked circular DNAs of two kinds. The maxicircles encode 9S and 12S rRNAs, two guide RNAs (gRNAs), and 18 proteins. The minicircles bear gRNA genes. Both maxicircle and minicircle genomes are transcribed from multiple promoters into 3' extended precursors which undergo 3'-5' exonucleolytic trimming. For translation, most pre-mRNAs further proceed through 3' adenylation, and often undergo U-insertion/deletion editing, and 3' A/U-tailing. The rRNAs and gRNAs are typically 3'

Highlights

- Mitochondrial RNA processing events in kinetoplastid protists include 5' modifications, 3'-5' degradation, internal sequence changes by U-insertion/deletion mRNA editing, and non-templated 3' extensions.
- The specificity of mRNA editing is dictated by guide RNAs while 5' modifications and 3' extensions are controlled by diverse PPR RNA binding factors.
- Antisense transcription plays a central role in delimiting 3'-5' trimming of primary transcripts.
- Macromolecular protein and ribonucleoprotein complexes and auxiliary factors involved in these processes have been identified and characterized to varying degrees. This review discusses recent developments and introduces a consensus nomenclature for mitochondrial RNA processing complexes and factors in *T. brucei*.

Outstanding Questions

- Recent understanding that mitochondrial mRNA, gRNA and rRNA genes are transcribed individually as 3' extended precursors places the onus on timing and mutual dependence of RNA synthesis and processing events. The nature of mitochondrial promoters and terminators, the composition of the transcription complex, the functionality of its components and the precise role of antisense transcripts remain to be elucidated.
- Many RESC subunits lack recognizable motifs and yet several bind RNA and most are essential for editing and cell viability. Understanding their functions would be a feast for structural biology.
- The mitochondrial ribosome likely selects fully-edited A/U-tailed mRNAs, but the mechanism of translation initiation at the 5' end is unclear.
- Mitochondrial proteomics remain a challenge due to extremely hydrophobic nature of mitochondrially-encoded proteins.
- The stage-specific patterns of mitochondrial transcription, RNA processing and translation are likely coordinated with nuclear gene expression by yet unknown mechanisms.

Lexis and Grammar of Mitochondrial RNA Processing in Trypanosomes

**Inna Aphasizheva,^{1*} Juan Alfonzo,² Jason Carnes,³ Igor Cestari,⁴ Jorge Cruz-Reyes,⁵
H. Ulrich Göringer,⁶ Stephen Hajduk,⁷ Julius Lukeš,⁸ Susan Madison-Antenucci,⁹
Dmitri A. Maslov,¹⁰ Suzanne M. McDermott,³ Torsten Ochsenreiter,¹¹ Laurie K.
Read,¹² Reza Salavati,⁴ Achim Schnauffer,¹³ André Schneider,¹⁴ Larry Simpson,¹⁵
Kenneth Stuart,³ Vyacheslav Yurchenko,¹⁶ Z. Hong Zhou,¹⁵ Alena Zíková,⁸ Liye
Zhang,¹⁷ Sara Zimmer¹⁸ and Ruslan Aphasizhev¹**

¹Department of Molecular and Cell Biology, Boston University Medical Campus, Boston,
MA 02118, USA

²Department of Microbiology, The Ohio State University, Columbus, OH 43210, USA

³Center for Global Infectious Disease Research, Seattle Children's Research Institute, Seattle,
WA, USA

⁴Institute of Parasitology, McGill University, Ste-Anne-de-Bellevue, Québec, Canada

⁵Department of Biochemistry and Biophysics, Texas A&M University, College Station, TX
77843, USA

⁶Department of Molecular Genetics, Darmstadt University of Technology, 64287 Darmstadt,
Germany

⁷Department of Biochemistry & Molecular Biology, University of Georgia, Athens, GA
30602, USA

23 ⁸Institute of Parasitology, Biology Centre, Czech Academy of Sciences and Faculty of
 24 Sciences, University of South Bohemia, České Budějovice (Budweis), Czech Republic

25 ⁹Parasitology Laboratory, Wadsworth Center, New York State Department of Health, Albany
 26 NY 12201, USA

27 ¹⁰Department of Molecular, Cell, and Systems Biology, University of California - Riverside,
 28 Riverside, CA 92521, USA

29 ¹¹Institute of Cell Biology, University of Bern, Switzerland

30 ¹²Department of Microbiology & Immunology, University at Buffalo, Jacobs School of
 31 Medicine and Biomedical Sciences, Buffalo, NY 14203, USA

32 ¹³Institute of Immunology and Infection Research, University of Edinburgh, Edinburgh EH9
 33 2FL, United Kingdom

34 ¹⁴Department of Chemistry and Biochemistry, University of Bern, Bern CH-3012,
 35 Switzerland

36 ¹⁵Department of Microbiology, Immunology and Molecular Genetics, University of
 37 California, Los Angeles, United States

38 ¹⁶Life Science Research Centre, Faculty of Science, University of Ostrava, Ostrava, Czech
 39 Republic; Martsinovsky Institute of Medical Parasitology, Sechenov University, Moscow,
 40 Russia

41 ¹⁷School of Life Science, Shanghai Tech University, Shanghai 201210, China

42 ¹⁸University of Minnesota Medical School, Duluth campus, Duluth, MN 55812 USA

43

44 ***Correspondence:** innaaf@bu.edu (I. Aphasizheva)

45 **Keywords:** *Trypanosoma*; mitochondria; kinetoplast; RNA editing; RNA decay;
46 polyadenylation; translation.

47

Abstract

Parasitic protist *Trypanosoma brucei* sp. cause African human and animal trypanosomiasis, a spectrum of diseases affecting the population and economy in sub-Saharan Africa. These dioxenous hemoflagellates belong to the class Kinetoplastea, a taxon distinguished by the presence of a kinetoplast. This nucleoprotein body contains mitochondrial DNA of two kinds: maxicircles and minicircles. The maxicircles (~25 per genome, each ~23 kb) encode 9S and 12S ribosomal RNAs, two guide RNAs (gRNAs), two ribosomal proteins and 16 subunits of respiratory complexes. The minicircles (~5,000 per genome, ~1 kb each) bear gRNA and gRNA-like small RNA genes. Relaxed maxicircles and minicircles are interlinked and packed into a dense disc-shaped network by association with histone-like proteins. Both maxicircle and minicircle genomes are transcribed by a T3/T7 phage-like RNA polymerase from multiple promoters into 3' extended precursors which undergo 3'-5' exonucleolytic trimming. To function in mitochondrial translation, most pre-mRNAs must further proceed through 3' adenylation, and often undergo gRNA-directed uridine insertion/deletion editing, and 3' A/U-tailing. The ribosomal and guide RNAs are typically 3' uridylated. Historically, the fascinating phenomenon of RNA editing has attracted major research effort, and recent developments provided insights into pre- and post-editing processing events and identified key players in transforming primary transcripts into mature RNAs, and regulating their function and turnover. Here, we introduce a unified nomenclature of mitochondrial RNA processing complexes and factors in *T. brucei* and attempt to integrate known modalities of kinetoplast RNA metabolism.

Mitochondrial Gene Expression in Trypanosomes: A Trove of Unconventional Biology

Protist parasites of the genus *Trypanosoma* have occupied the research spotlight since 1895 when David Bruce identified *Trypanosoma brucei* as the causative agent of animal trypanosomiasis (Nagana) [1], and later works linked these organisms to sleeping sickness in humans [2]. Biomedical, economic and societal impact of parasite infections warranted in-depth studies of the fascinating biology underlying *T. brucei* metabolism, development and interactions with insect vector and mammalian hosts [3]. Among the most striking cellular features of these parasites is the bipartite mitochondrial genome consisting of maxicircles and minicircles, and aggregately referred to as the kinetoplast DNA (kDNA). In *T. brucei*, maxicircles are catenated with minicircles into a single network and compacted by histone-like basic proteins. The progress in elucidating kDNA maintenance and segregation has been reviewed elsewhere [4-6]. Maxicircles, an equivalent of mitochondrial genomes (mtDNA) in other organisms, encode 9S and 12S rRNAs, two ribosomal proteins [7], and 16 subunits of respiratory complexes. Unlike most organellar genomes, kDNA lacks tRNA genes [8, 9], and 12 maxicircle genes are present as cryptogenes whose transcripts require RNA editing to restore a protein-coding capacity [10]. The editing is mediated by hundreds of guide RNAs (gRNAs) which are mostly encoded by minicircles, with only two gRNAs encoded by maxicircles. The distinct, albeit interlinked, maxicircle and minicircle genomes are transcribed independently, but the information converges at the post-transcriptional level whereby minicircle-encoded gRNAs direct editing of maxicircle encoded pre-mRNAs. The evolution of editing and whether this process confers a selective advantage to kinetoplastids remain the subject of a debate [11], but the existence of alternatively edited mRNA sequences and cognate gRNAs raises a possibility that editing-driven protein diversity may be functionally relevant [12-15]. Historically, much attention has been focused on RNA editing

mechanism and composition of editing complexes [16-22] while more recently major advances have been made in understanding transcription [23], primary RNA nucleolytic processing [24-26], 5' [23] and 3' modifications [27-32], and ribosome biogenesis and translation processes [7, 33-35]. The perceived complexity of mitochondrial gene expression has been exacerbated by recent influx of new RNA processing factors and numerous names often referring to the same entity. Bearing in mind that functions of only few proteins and complexes are established beyond reasonable doubt, we nonetheless submit that the process of discovering major players is close to completion. Here, we outline major stages in kinetoplast RNA processing and build on previous attempts [36, 37] to introduce a unified nomenclature for respective protein and ribonucleoprotein complexes, enzymes and factors in *T. brucei*. Given that orthologs of nearly all *T. brucei* proteins listed in Table 1 are readily distinguishable in related organisms, this nomenclature should be broadly applicable to other *Kinetoplastea*.

Nucleolytic Processing of Primary Transcripts

In most organisms, primary polycistronic mtDNA transcripts are punctuated by tRNAs whose excision by RNases P and Z defines functional RNA boundaries [38]. Although loss of mtDNA-encoded tRNA genes renders such a mechanism inapplicable to *T. brucei*, it has been held that mature mRNAs and rRNAs with uniformly monophosphorylated 5' and well-demarcated 3' termini reflect maxicircle polycistronic precursor's partitioning by an unknown endonuclease [39, 40]. To that end, a prominent transcription start site has been mapped within the divergent region ~1,200 nt upstream of 12S rRNA [41] and transcription proceeding through intergenic regions has been reported [42]. The uridylylated rRNA [43] and adenylated mRNA [44] termini also typify distinct 3' end modification mechanisms for these

RNA classes. Conversely, short (30-60 nt) gRNAs maintain 5' triphosphates characteristic of transcription-incorporated initiating nucleoside triphosphate and, similar to rRNAs, are 3' uridylated [45]. However, the only explored candidate gRNA precursor processing endonuclease KRPN1 (mRPN1) [46] is dispensable for axenically grown bloodstream stage parasites [42]. Thus, it may be argued the essentiality of RNA editing [47], which requires mature gRNAs [25, 48], renders KRPN1 an unlikely contributor to gRNA precursor processing. The observations conducive to the endonucleolytic model have been re-examined in light of mitochondrial 3' processome (MPsome) discovery [24]. These studies recognized the MPsome-catalyzed 3'-5' exonucleolytic degradation as the major nucleolytic processing pathway for mRNA and rRNA [29], and for gRNA [24] precursors. Composed of KRET1 TUTase [49], KDSS1 exonuclease [26, 50] and mitochondrial processome subunits (MPSS) 1-6 (Table 1), the purified MPsome displays 3'-5' RNA degradation, RNA hydrolysis-driven double-stranded RNA unwinding, and 3' RNA uridylation activities. Although the autonomous KDSS1 is inactive, incorporation into the MPsome converts this polypeptide into a highly processive exonuclease capable of degrading structured RNAs to 5-6 nt fragments. On the other hand, individual KRET1's robust UTP polymerization activity [51] is tamed upon MPsome assembly to adding 1-15 Us, a pattern consistent with the U-tails observed in steady-state RNA [15, 52]. Cumulatively, detection of *in vivo* uridylated precursors and degradation intermediates [24, 31], stimulation of *in organello* KRET1-dependent RNA decay by UTP [27], and MPsome's preference for U-tailed substrates suggest that uridylation by KRET1 activates RNA degradation by KDSS1. It is unclear whether substrate tunneling occurs within the same particle, but a coupling between RNA uridylation and degradation by 3'-5' RNase II/RNB-type exonuclease appears to be a highly conserved and phylogenetically widespread mode of RNA decay [53, 54].

Exonucleolytic processing is often a case of regulated decay whereby mature 3' termini are defined by a degradation blockade at a specific sequence or structure. In the *T. brucei* mitochondrion, antisense transcripts cause MPsome pausing at 10-12 nt before double-stranded region at which point MPsome-embedded KRET1 likely adds a U-tail causing disengagement from the RNA [23, 24]. It follows that the precise transcription initiation site on the antisense strand defines the position of mature 3' terminus of the sense transcript. The antisense model of gRNA 3' end definition is consistent with bi-directional transcription from converging promoters otherwise recognized as imperfect 18-nt inverted repeats [55] that flank almost all gRNA genes in *T. brucei* minicircles [56, 57]. Identification of gRNA-sized short antisense RNAs and accumulation of antisense precursors upon KRET1 and KDSS1 knockdowns [24] further indicates that sense and antisense precursors hybridize with their complementary 5' regions. In the current model, the length of a double-stranded region, which is a distance between respective transcription start sites, likely defines gRNA length prior to uridylation [24]. However, most *T. brucei* minicircles encode 3-4 gRNA cassettes [57] and primary RNAs may exceed a 1 kb linear length of a minicircle [24, 25]. Hence, an extensive transcription of both strands may also generate much longer double-stranded RNAs that are degraded by yet unknown mechanism. Conversely, *L. tarentolae* minicircle typically contains a single gRNA gene and lacks recognizable inverted repeats [58]. Although both strands are transcribed [59], the gRNA-flanking sequences are dissimilar from those of *T. brucei*, which indicates a divergent nature of minicircle promoters among trypanosomatids. The maxicircle promoters remain to be determined, but detection of antisense transcription start sites near intergenic regions and presence of corresponding non-coding antisense transcripts make a reasonable argument for a general mechanism of 3' end definition for minicircle and maxicircle transcripts [23].

Modification of the 5' End

The 5' monophosphorylated termini of maxicircle-encoded rRNAs and mRNAs have long been interpreted as indicative of endonucleolytic partitioning of polycistronic precursors. It is, however, unfeasible to produce more than one monocistronic mRNA from a precursor by 3'-5' degradation. This logic dictates that: 1) each gene rests under control of a dedicated promoter; 2) the 5' terminus is set by transcription initiation; 3) inorganic pyrophosphate (PPi) is selectively removed from initiating nucleoside triphosphate in mRNAs and rRNAs, but not in gRNAs; and 4) transcription may proceed across multiple genes and produce a 3' extended precursor of which only the most 5' coding region is preserved after 3'-5' trimming. Identification of the 5' pyrophosphate processome complex (PPsome) partially resolved the question of differential phosphorylation status and linked 5' PPi removal to mRNA stability [23]. A stable protein complex of MERS1 NUDIX [Nucleoside diphosphate linked to (X)] hydrolase and MERS2 PPR RNA binding subunit, the MPsome selectively binds to degenerate G-rich motifs found near mRNA 5' ends, but not in gRNAs. MERS1 hydrolase is catalytically inactive as an individual protein while MERS2 confers both binding specificity and affinity for RNA substrate. Remarkably, MERS1 downregulation or replacement with an inactive version effectively eliminates most mRNAs but exerts negligible effects on gRNAs and rRNAs. It appears that rRNA is stabilized by different factors, possibly those involved in ribosome biogenesis [35]. Although PPsome-dependent mRNA protection against 3'-5' degradation (see below) and the essential role of PPi removal are evident, the mechanistic insights into these processes will likely come from understanding PPsome's interactions with RNA editing substrate binding (RESC) and polyadenylation (KPAC) complexes discussed below [32].

Modifications of the 3' end

Non-templated 3' nucleotide additions often wield profound influence on RNA processing, function, trafficking and turnover [60]. In *T. brucei*, mitochondrial RNA 3' modifications are classified into U-tailing by KRET1 TUTase (gRNAs and rRNAs), A-tailing by KPAP1 poly(A) polymerase (most mRNAs [31]), and A/U-extensions which require both enzymes and a complex of Kinetoplast Polyadenylation Factors 1 and 2 (KPAF1/2, [30]). Lack of pronounced RNA substrate specificity for KRET1 and KPAP1 raises the question of accessory factors that enable modifications of distinct RNA classes, and the functionality of these extensions. The presence of short U-tails in gRNAs and rRNAs, as well as non-templated uridine residues sometimes found in mRNAs between the 3' UTR and the A-tail [30], indicates that uridylation by the MPsome-embedded KRET1 is a default 3' modification. It is plausible that U-tailing causes the MPsome to disengage from the precursor when degradation pauses near double-stranded region formed by antisense RNA. However, the U-tail itself does not exert an appreciable impact on mature gRNA or rRNA stability [25] and its functionality beyond termination of processing remains debatable [61-63]. Conversely, a short (15-30 nt) A-tail decorates most mRNAs and impacts stability depending on transcript's editing status [28, 31, 64]. As demonstrated by KPAP1 poly(A) polymerase loss-of-function studies [29, 31] and *in organello* decay assays [28], adenylation mildly de-stabilizes pre-edited transcripts only to become essential for maintaining RNAs that are edited beyond initial editing sites at the 3' end. A short A-tail also stabilizes never-edited mRNAs (those that contain an encoded open reading frame and do not require editing). The coupling between an mRNA's editing status and opposing effects of adenylation points toward a surveillance system capable of both sensing the extent of internal U-insertions/deletions and enabling 3' A-tail addition and function. In molecular terms, sequence-specific activators and inhibitors would be expected to modulate mRNA

adenylation by KPAP1, and the resistance of such a modified molecules to decay. The respective functions have been attributed to Kinetoplast Polyadenylation Factors (KPAF) 3 [29] and 4 [32], which belong to a family of 35 amino acid repeat-containing RNA binding (PPR) proteins. Discovered in land plants [65], the helix-turn-helix PPR motif recognizes a single nucleoside via side chains occupying cardinal positions 5 and 35 of the repeat (or the last position in a longer structure). An array of adjacent PPR repeats often folds into a superhelical domain capable of binding to specific RNA sequence and recruiting or blocking various enzymes [66-68]. In this context, KPAF3 reportedly binds to G-rich pre-edited mRNAs with sufficient affinity and coverage to stabilize these species following 3'-5' trimming by the MPsome [29]. *In vitro* reconstitution experiments further demonstrate that KPAF3 stimulates KPAP1 polyadenylation activity and this effect depends on the presence of the G-rich site near the 3' end. Remarkably, KPAF3 binding is eliminated by the initiating editing events leaving the stability of edited RNA reliant on A-tail added prior to editing [29]. Thus, KPAF3 functions as editing sensor and *bona fide* polyadenylation factor thereby connecting the internal sequence changes and 3' modification [29-31].

The most apparent A-tail function would be protecting mRNA against degradation by the MPsome. However, *in vitro* studies show that A-tailed RNAs can be degraded by the purified MPsome, albeit less efficiently than uridylylated substrates [24]. The A-tailed partially-edited pre-mRNAs are also somehow prevented from the post-editing addition of the 200-300 nt-long A/U-tail. This modification marks fully-edited molecules [31] and channels translationally-competent mRNA for translation [30, 33, 34]. Finally, the mechanism of mRNA stabilization by the PPsome must reconcile binding of this complex to the 5' end with blocking 3'-5' degradation [23]. To rationalize these observations, Mesitov et al., envisaged a *trans*-acting factor that recognizes a nascent A-tail to enable an interaction between PPsome occupying the 5' end and polyadenylation complex (KPAC) bound to the 3'

end [32]. It has been proposed that the resultant circularization increases mRNA resistance to degradation and uridylation, and, therefore, to premature A/U-tailing and translational activation of partially-edited transcripts [32]. Trypanosomal genomes apparently lack mitochondrially-targeted canonical RRM motif-containing poly(A) binding proteins, but this function is fulfilled by KPAF4. This PPR protein is almost entirely composed of seven repeats of which five are predicted to bind sequential adenosine residues [69]. Co-purification studies support KPAF4 interactions with KPAC components (KPAP1, KPAF1/2) and RESC-mediated contacts with the PPsome. Accordingly, the A-tail has been identified as the predominant *in vivo* binding site while *in vitro* KPAF4 selectively recognizes adenylated substrates. Indeed, in the presence of KPAF4 adenylated RNA is more resistant to degradation by the purified MPsome and uridylation by KRET1 TUTase.

Although direct demonstration of mRNA circularization is lacking, this event can be imagined as a critical quality check point to ensure 5' end occupancy by the PPsome and correct termination of 3'-5' trimming downstream from the KPAF3 binding site. In this scenario, KPAF3 binding likely selects a correct 3' UTR among trimmed precursor isoforms thereby distinguishing mRNA from rRNA and stimulating polyadenylation of the former by KPAP1. KPAF4 binding to a nascent A-tail would enable interaction with 5' end-bound PPsome, hence, mRNA circularization. Consequentially, only A-tailed mRNAs would proceed through the editing cascade while the variants truncated beyond KPAF3 binding sites become uridylated and degraded [29]. It follows that upon editing completion at the 5' end, a signaling event must take place to release the circularization and enable access of KPAF1/2 factors and KRET1 TUTase in order to add long A/U-tail to the pre-existing short A-tail. Although these postulates require further testing, it seems plausible that PPsome displacement from the 5' end by final editing events may disrupt circularization and stimulate A/U-tailing, a final processing step that renders mRNA competent for translation.

U-insertion/deletion mRNA Editing

Editing Process

In *T. brucei*, six of the 18 annotated mRNAs encode predicted polypeptides while the remaining 12 transcripts undergo editing to acquire a protein-coding sequence. The extent of editing varies from minor, typified by insertion of four Us into three closely-spaced sites (COII mRNA, [10]), to moderate (e.g., *cyb* mRNA, 34 Us are inserted into a confined region near 5' end [70]), to pan-editing during which hundreds of uridines are inserted or deleted throughout the entire transcript (e.g. ND7 mRNA, [71]). The determinants of position-specific U-insertions and deletions were discovered in the Simpson laboratory as short patches of complementarity between edited mRNA and maxicircle DNA in *Leishmania tarentolae* [45]. By allowing for G-U, in addition to canonical Watson–Crick base-pairing, short (50-60 nt) RNAs transcribed from minicircles have been recognized as carriers of genetic information and termed guide RNAs (gRNAs). *In vitro* experiments by the Stuart laboratory directly demonstrated that gRNAs indeed constitute the necessary and sufficient source of editing information [72-74]. Although the first gRNAs were discovered in the maxicircle, further work in *Leishmania* established that most gRNAs are encoded in minicircles [75]. In *T. brucei*, only two gRNAs have been identified in the maxicircle: a *cis*-acting element embedded into 3' UTR of COII mRNA [76], and a *trans*-acting gRNA that completes editing of the MURF2 mRNA. The secondary structure of gRNA-mRNA dictates the editing site selection and the extent of U-insertions and deletions [72]. The initial gRNA-mRNA interaction is accomplished via a short (10-12 nt) region of complementarity between the gRNA's 5' anchor region and the pre-edited mRNA. The remaining guiding segment forms an imperfect duplex with pre-mRNA resulting in looping out of single-stranded uridines in mRNA (deletion sites) or purine nucleotides in gRNA (insertion sites). At either site, the mRNA is cleaved at the first unpaired nucleotide adjacent to the 5' anchor duplex.

The resultant structures of deletion and insertion intermediates are distinct: single-stranded uridines become exposed to a 3'-5' exonucleolytic attack in the former, while a single-stranded gap is created between two helices in the latter. Upon trimming single-stranded uridines from the 5' cleavage fragment in the deletion site or adding gap-specified number of Us into the insertion site, the fragments are joined to restore mRNA continuity. Both types of sequence changes extend the double-stranded anchor region. Pan-editing requires multiple overlapping gRNAs and there is a method to it: sequence changes directed by the initiating gRNA create a binding site for the next one to ensure an overall 3'-5' polarity along the editing domain. However, editing may not always proceed strictly 3' to 5' as "mis-edited" junctions are present at the leading edge of editing in the majority of steady-state mRNAs [59, 77-79]. The role of junctions is not entirely understood, but they likely represent a mixture of regions that undergo re-editing to canonical edited sequence, dead-end products, or mRNAs with alternative non-canonical coding sequences [22].

A single editing domain may cover an entire mRNA [80], or an isolated region [71]. An individual gRNA can theoretically direct insertions and deletion at several closely spaced sites (editing block), but as editing progresses within the block, the interaction between gRNA and mRNA 5' cleavage fragment is supported by fewer base pairs. Stabilizing the 5' cleavage fragment-mRNA tethering by additional base pairing stimulates cleavage and the full editing cycle *in vitro* [81-83], but it is unclear how the problem of editing at distal sites within one block or across sequential blocks is solved *in vivo*. An active displacement of a gRNA with diminishing "3' tether" by RNA helicase is among possible solutions that would enable binding of succeeding gRNA within a domain. Alternatively, the gRNA's U-tail may help stabilize interaction with an mRNA [61, 84], but the conclusive evidence for this event is yet to be obtained.

Elemental Editing Reactions

Editing reactions are catalyzed by enzymes embedded into the ~20S (~800 kDa) RNA Editing Catalytic Complex (RECC), a remarkable example of a modular assembly that enables broad functionality on distinct RNA substrates [85-89]. A common core particle consists of U-insertion (KRET2 TUTase, KREPA1 zinc finger protein and KREL2 RNA ligase) and U-deletion (KREX2 exonuclease, KREPA2 zinc finger protein and KREL1 RNA ligase) sub-complexes and six structural and/or RNA binding proteins (KREPA3, KREPA4, KREPA5, KREPA6, KREPB4, KREPB5). The U-insertion and U-deletion sub-complexes likely function independently [48, 90] while most of the remaining components are essential for assembly and/or integrity of the entire core particle [91-94]. The core particle is shared among at least three RECC isoforms distinguished by association with endonuclease modules. Each module is composed of an RNase III endonuclease and a partner protein(s) and is primarily responsible for recognition and cleavage of insertion and deletion sites. The U-deletion sites are recognized by the RECC isoform with KREN1+KREPB8+KREX1, while U-insertion sites are recognized by the RECC isoforms with KREN2+KREPB7 or KREN3+KREPB6, which display distinct and overlapping specificities [95-99]. The canonical RNase III catalytic domain typically forms a functional homodimer with two active sites that introduce four cuts into both strands of a double-stranded RNA [100]. In contrast, editing endonucleases appear to cleave only mRNA. It seems plausible that RNA hydrolysis is restricted to a single cut by heterodimer formation between KREN1, KREN2 or KREN3, and catalytically-inactive degenerate RNase III domains in KREPB4 or KREPB5 [101]. A contribution of RNase III partner proteins KREPB8, KREPB7 or KREPB6 to modulating cleavage activity is also possible [102, 103]. Binding of KREN1, KREN2 and KREN3 modules to a common core containing U-deletion, U-insertion and ligase activities highlights RECC's modular nature [98, 104, 105], but the nature of interactions responsible for mutually

exclusive contacts between the core and distinct modules remains unclear. Cross-linking mass-spectrometry points to interactions involving RNase III domain dimerization between editing endonucleases with partner proteins KREPB6, B7 or B8, and core proteins KREPB4 and KREPB5 [102, 103, 106].

Within the common core, the U-deletion and U-insertion cascades are spatially separated by virtue of editing enzymes binding to zinc finger proteins, KREPA2 and KREPA1, respectively [90, 107-109]. KREX1 and KREX2 proteins possess exonuclease-endonuclease-phosphatase (EEP) catalytic domains and display single-stranded uridine-specific 3'-5' exonuclease activity *in vitro* [108, 110]. However, their protein-protein interactions are remarkably distinctive: the essential KREX1 belongs to the KREN1 endonuclease module, and is responsible for the main U-deletion activity; the dispensable KREX2 probably represents a structural component of the U-deletion sub-complex [96, 106]. Fittingly, *L. tarentolae* KREX2 lacks a catalytic domain, but remains associated with the U-deletion sub-complex [85]. In the U-insertion sub-complex, KRET2 TUTase binds to KREPA1, which results in a mutual stabilization and stimulation of TUTase activity [48, 106, 111-113]. Selectivity of uridine incorporation is determined by KRET2's intrinsic specificity for UTP [114] rather than the nature of the opposing nucleotide in the gRNA. To that end, the +1U addition occurs equally efficient irrespective of the corresponding nucleotide in gRNA, but the +2U addition occurs only if the +1U forms a base pair with either adenosine or guanine. Consequentially, both purine bases in guiding positions direct U-insertions with similar efficiency [81, 111]. RNA editing ligases 1 and 2 (KREL1 and KREL2) have been identified as components of U-deletion and U-insertion sub-complexes, respectively [90, 106, 107]. Although spatial separation appears to suggest specialized roles, only KREL1, but not KREL2, is essential for cell viability [47, 115, 116].

Editosome Definition

From the early reports of RNA ligase-containing complexes sedimenting in glycerol density gradients as particles with apparent 20S to 50S values [117, 118], the quest for an elusive “editosome” evolved into a concept of an RNA editing holoenzyme. For the purposes of this review, we shall equate the editosome and editing holoenzyme and define this entity as an RNA-mediated assembly of the RNA editing catalytic (RECC), RNA editing substrate binding (RESC) and RNA editing helicase REH2 (REH2C) complexes. It is a virtual certainty that additional components are also involved [19, 21, 119]. This definition stems from parallel lines of inquiry by the Stuart, Afasizhev and Lukeš laboratories that identified an ~800 kDa protein complex (originally termed Mitochondrial RNA Binding Complex 1, MRB1, and Guide RNA Binding Complex, GRBC), of which two components are essential for gRNA stability [120-122]. Initially named GRBC1 and GRBC2, these homologous polypeptides lack annotated motifs and similarity to any protein outside of Kinetoplastids [63]. GRBC1 and GRBC2, also referred to as GAP2 and GAP1, respectively [122], form a stable heterotetramer which binds gRNA *in vitro* and *in vivo* [52]. Extensive co-purification and yeast two-hybrid screens further dissected MRB1 into two relatively stable protein complexes: an ~20-component RNA Editing Substrate Binding Complex (RESC), which includes RESC1 (GRBC1, GAP2) and RESC2 (GRBC2, GAP1), and three-subunit RNA Editing Helicase 2 Complex (REH2C) (Table 1). It appears that both RESC and REH2C bind editing substrates, intermediates and products, and engage in RNA-mediated interactions with the catalytic RECC complex [52, 123-126]. All but five of the RESC subunits lack discernible motifs or similarities to non-kinetoplastid proteins, although several exhibit *in vitro* RNA binding activity [19, 21, 119]. Most subunits are essential for cell viability and their knockdowns typically produce phenotypes consistent with an inhibited editing process. Recently, a substantial progress has been made in deciphering roles of individual factors. The

RESC1/2 tetramer appears to be solely responsible for gRNA stabilization [121, 122]. Deep sequencing studies showed that the RRM/RGG-containing RESC13 (RGG2) and proximal protein RESC11A (MRB8180) contribute to editing processivity within an extended domain [78, 127]. These two proteins strongly promote the formation of junctions, implying a critical role of these regions in editing progression [78]. Conversely, the product of a duplicate gene RESC12A (MRB8170) has been implicated in editing initiation and in constraining the region of active editing [78, 128]. Biochemical attempts to refine RESC architecture indicate a modular organization with potential protein clusters responsible for interaction with the RECC and polyadenylation complexes [52, 129]. However, an unambiguous assignment of specific polypeptides to functionally meaningful modules awaits elucidation of a high-resolution structure of the RESC complex, which is likely to be heterogenous and dynamic.

RNA editing is an essential processing step for a subset of mitochondrial transcripts and must be integrated into a general pathway of producing translation-competent mRNAs. Accumulating evidence suggests that the RESC is responsible for coordinating pre- and post-editing processing events via RNA-mediated contacts with 5' and 3' modification complexes, and auxiliary factors. On the other hand, the catalytic RECC isoforms appear to act on RESC-bound editing substrates in a transient manner. The RNA-mediated interaction between RESC and PPsome has been deduced from co-purification of RESC1/2 (GRBC1/2) and MERS1 hydrolase [23, 121, 130] whereas *in vivo* proximity biotinylation identified RESC19 (MERS3) as the most plausible adapter subunit [23]. An independent study predicted Z-DNA-binding domains in RESC19 (termed RBP7910) and showed that *in vitro* this protein preferentially binds RNAs containing poly(U) and poly(A/U)-rich sequences [131]. Likewise, kinetoplast polyadenylation complex components have been consistently detected in various RESC preparations and particularly in those with tagged RESC15-18 [29, 32, 52]. Furthermore, RESC13 (RGG2) and surrounding proteins likely mediate contacts between

RESC and RECC complexes [52]. Finally, RNA editing helicase 2 (KREH2, see below) preferentially associates with RESC variants purified by tagging of either RESC1 or RESC2 [52, 121, 132-134]. Thus, the RESC complex functions not only in binding of RNA editing substrates, intermediates and products, but also recruits mRNA modification complexes and auxiliary factors via specific subunits.

The Kinetoplast RNA Editing Helicase 2 (KREH2) complex, termed REH2C, consists of DEAH/RHA RNA helicase KREH2, and KREH2-associated factors 1 (KH2F1) and 2 (KH2F2). KH2F1 contains eight C2H2 zinc fingers while KH2F2 lacks any identifiable motifs [135] (Table 1). Isolated REH2C exhibits ATP-dependent 3'-5' dsRNA unwinding activity and co-sediments with a major peak of same activity in mitochondrial extracts [134]. Zinc finger protein KH2F1 emerged as an adaptor connecting KREH2 helicase with the editosome while gRNA-mRNA hybridization has also been implicated in facilitating this interaction [135]. KREH2 and KH2F1 knockdowns display consistent phenotypes of increased editing pausing and reduced processivity of editing, which are indicative of REH2C participation in continuous remodeling of gRNA-mRNA contacts and perhaps the entire editosome [123, 135]. It must be emphasized that the reciprocal affinity purifications remain the most salient evidence of the editosome being an RNA-based assembly of RECC, RESC and REH2C protein complexes [120-122, 133, 134].

Auxiliary RNA Processing Factors

Putative Poly(A) Polymerase KPAP2

A putative kinetoplast poly(A) polymerase KPAP2 has been identified by homology to the human mitochondrial enzyme and apparently is not required for axenic *T. brucei* growth in either bloodstream or procyclic life stages [136]. Although KPAP2 protein sequence is highly

similar to that of KPAP1, its enzymatic identity and *in vivo* substrates remain to be established. Available proteomics data do not support KPAP2 association with KPAC [29, 31, 32].

REH1 RNA Helicase

Editing reactions are expected to produce an mRNA-gRNA duplex wherein gRNA must be eventually displaced to allow binding of a sequential gRNA, or before the edited transcript can be translated. It stands to reason that active remodelers, such as DEAD/H-box RNA helicases, would be involved and indeed two such proteins have been implicated in the editing process. However, demonstrating their specific roles, RNA targets and mechanism of action proved to be challenging. Knockdown of KREH1 (Hel61) helicase [137, 138] affected editing mediated by two or more overlapping gRNAs [139], but the existing data do not allow unambiguous placing of KREH1 into mRNA editing or processing complexes. The nature of KREH1 substrates and the timing and purpose of KREH1-dependent RNA remodeling also remain unclear. Although much remains to be understood about the helicases' involvement in editing and other processes, the transcript-specific impacts of repressing KREH1 and KREH2 (discussed above) justify further efforts to dissect their molecular functions.

RNA Binding Factors

Kinetoplast Mitochondrial RNA Binding Proteins 1 and 2 (KMRP1 and 2), originally called gBP21 and gBR25, then MRP1 and MRP2, were identified independently in *T. brucei* by UV-induced crosslinking with synthetic gRNA (gBP21, [140]), in *C. fasciculata* as poly(U) binding proteins (gBP21 and gBP25, [141]), and in *L. tarentolae* via crosslinking to double-stranded RNA resembling the U-deletion site (MRP1 and MRP2, [130]). Extensive biochemical and structural studies concluded that KMRP1 and KMRP2 assemble into an

~100 kDa heterotetramer, which binds both single- and double-stranded RNAs with high affinity [130, 142, 143]. These RNA binding properties are manifested by an RNA annealing activity, an attractive accessory function that may promote gRNA binding to cognate mRNA targets [144, 145]. However, the transcript-specific impact of dual KMRP1/2 repression suggests a contribution to stabilization of moderately edited and some never-edited mRNAs rather than direct participation in the editing process [142, 146, 147]. In support of this notion, RNAi experiments demonstrated that MRP1/2 depletion virtually eliminates the edited form of the moderately-edited *cyb* mRNA, but exerts little impact on the pre-edited transcript [146-148]. While much is known about the KMRP1/2 structure and *in vitro* properties, the definitive function of this RNA binding complex remains to be established. Much of the same narrative applies to KRBP16 (RBP16), which carries N-terminal cold shock and C-terminal RG-rich domains [149]. RNAi studies revealed an overlap between mRNA sets negatively affected by individual KMRP1/2 and KRBP16 knockdowns: edited *cyb* mRNA, but not any other edited transcripts, was severely downregulated while never-edited CO1 and ND4 transcripts also declined [146]. KRBP16 *in vitro* properties, such as RNA binding affinity, RNA annealing activity and stimulation of editing activity [146, 148-155], and the impact of RNAi knockdown on the initiation of *cyb* mRNA editing [148] are consistent with participation in the editing process, although the mechanistic role remains to be firmly established. Another enigmatic RNA binding protein, KRGG1 was serendipitously discovered in a large (>50S) RNP of unknown nature [156], and subsequently demonstrated to associate with the RESC complex [122, 157]. A different study identified a ribosome-bound KRGG1 fraction, which would explain the observed sedimentation patterns, but found no impact on RNA editing [52]. Another arginine-glycine-rich protein KRGG3, originally identified by association with RESC1/2 proteins and termed MRB1820 [129], is essential for parasite viability [158]. However, most KRGG3's interactions appear to be RNA-mediated

while the RNAi knockdown does not significantly impact major mitochondrial RNA classes. A structural study identified an ABC-ATPase fold and potential RNA binding surface in a 72 kDa protein KRBP72, initially termed MRB1590 [159]. KRBP72 knockdown specifically impacts editing of A6 mRNA [159]; however, an unequivocal functional placement of this factor also requires further investigation. Finally, participation of KREAP1 in editing [160, 161] has been contravened by a report of general mitochondrial RNA upregulation upon its knockdown and non-essentiality for parasite survival [162]. Mitochondrial RNA binding proteins are abundant and notoriously promiscuous in their interactions and pleiotropic effects on RNA steady-state levels [163], which makes an unequivocal definition of their function a challenge worth meeting.

Ribonucleases

Mitochondrial RNA processing most likely involves nucleolytic events beyond mRNA cleavage by editing KREN1, 2 and 3 endonucleases, KREX1 exonuclease, and 3'-5' degradation by the MPsome-embedded KDSS1. To that end, three distinct enzymes have been identified and characterized to various degrees. The single-strand uridine specific KRND1 3'-5' exonuclease [164] displays *in vitro* specificity for single-stranded uridines, similar to that of KREX1 editing enzyme [108, 110], and yet possesses an RND rather than EEP exonuclease domain. Given the diversity of U-tailed RNAs in the kinetoplastid mitochondrion, it is tempting to speculate on KRND1 involvement in regulating the 3' modification state, but its definitive function remains to be established. The same narrative applies to KRPN1, an RNase III-like endonuclease with a characteristic double-stranded RNA cleaving activity [42, 46] suggested to function in gRNA processing. Further studies are required to reconcile KRPN1 activity and interactions with an exonucleolytic mechanism of gRNA precursor processing by the MPsome. Finally, the discovery of PPR-repeat containing proteinaceous RNase P (PROPR2 [165], re-named here as KRNP1) supported earlier reports

of RNase P activity-like which removes the 5' leader from a synthetic tRNA precursor in mitochondrial lysate [166]. However, tRNAs are apparently imported into the mitochondrion with 5' and 3' extensions removed [167, 168], which leaves the nature of KRN1P1 *in vivo* substrates open to future inquiry.

Concluding Remarks

This review compiles 74 processing enzymes, RNA binding proteins and factors with unknown functionality that nonetheless are associated with RNA processing complexes. Proteomics and interactions analyses allowed clustering most of these into 5-6 macromolecular assemblies albeit with various degree of confidence. Although the list is almost certainly incomplete, with complexes and individual proteins, and their interactions and functions being constantly re-examined, the overall flow of RNA processing in trypanosome mitochondrion is taking shape and meaning. The key players responsible for maturation of 5' and 3' termini have been defined and initial insights gained into the molecular mechanism of internal sequence changes by editing. At this point, we suggest that the RNA editing holoenzyme (editosome) represents an RNP that chiefly includes three relatively stable protein complexes (RECC, RESC and REH2C) and RNA editing substrates, intermediates and products. It is understood that the definition of protein complex is to large extent a matter of purification techniques and we posit that future structural studies will shed light on stoichiometry and functions of individual subunits and modules. Because of sequence changes introduced by editing during mRNA processing, the overall picture is emerging of the mRNA fate being dictated by diverse PPR RNA surveillance factors. These proteins direct 5' pyrophosphate removal, transcript stabilization and pre-editing A-tailing, monitor initiation and progression of editing, and signal its completion by stimulating the

538 A/U-tailing. Displacement of bound PPRs from pre-edited mRNA by the editing process
539 appears to be the principal quality control mechanism. It remains to be established whether
540 active RNP remodeling takes place or sequence changes alone suffice for this purpose. In any
541 event, the plurality of PPRs and their capacity to read linear sequences and modulate the
542 activity of RNA modification and degradation enzymes position this protein family as the
543 focal point of mitochondrial RNA processing.

544

545

Acknowledgements

Afasizheva lab is supported by NIH grant AI113157. Afasizhev lab is supported by NIH grants AI091914 and AI101057. Cruz-Reyes lab is supported by NSF grant 1616845 and TAMU X-grant and T3 award. Göringer lab is supported by German Research Council (SPP1857). Lukeš lab is supported by the Czech Ministry of Education (OPVVV16_019/0000759 and ERC CZ LL1601. Read lab is supported by NIH grants AI061580, GM129041, and AI139448. Hajduk lab is supported by NIH grant AI125487. Schnauffer lab is supported by MRC Senior Non-Clinical Fellowship MR/L019701/1. Schneider lab is supported by the NCCR grant "RNA & Disease" and by the grant 175563 from the Swiss National Science Foundation. Stuart lab is supported by NIH grant 5R01AI014102-43. Yurchenko lab is supported by the European Regional Funds (project "Centre for Research of Pathogenicity and Virulence of Parasites", OPVVV16_019/0000759). Zíková lab is supported by the Czech Ministry of Education (OPVVV16_019/0000759) and by GACR grant 18-17529S. Zimmer lab is supported by NIH grant AI135885.

REFERENCES

1. Bruce, D. (1895) Preliminary report on the tsetse fly disease or nagana in Zululand, Bennett & Davis, printers.
2. Steverding, D. (2008) The history of African trypanosomiasis. *Parasit Vectors* 1 (1), 3.
3. Maslov, D.A. et al. (2019) Recent advances in trypanosomatid research: genome organization, expression, metabolism, taxonomy and evolution. *Parasitology* 146 (1), 1-27.
4. Schneider, A. and Ochsenreiter, T. (2018) Failure is not an option - mitochondrial genome segregation in trypanosomes. *J Cell Sci* 131 (18).
5. Povelones, M.L. (2014) Beyond replication: division and segregation of mitochondrial DNA in kinetoplastids. *Mol Biochem Parasitol* 196 (1), 53-60.
6. Jensen, R.E. and Englund, P.T. (2012) Network news: the replication of kinetoplast DNA. *Annu. Rev. Microbiol* 66, 473-491.
7. Ramrath, D.J.F. et al. (2018) Evolutionary shift toward protein-based architecture in trypanosomal mitochondrial ribosomes. *Science* 362 (6413).
8. Simpson, A.M. et al. (1989) Kinetoplastid mitochondria contain functional tRNAs which are encoded in nuclear DNA and also small minicircle and maxicircle transcripts of unknown function. *Nucleic Acids Res* 17, 5427-5445.
9. Hancock, K. and Hajduk, S.L. (1990) The mitochondrial tRNAs of *Trypanosoma brucei* are nuclear encoded. *J. Biol. Chem* 265, 19208-19215.
10. Benne, R. et al. (1986) Major transcript of the frameshifted coxII gene from trypanosome mitochondria contains four nucleotides that are not encoded in the DNA. *Cell* 46, 819-826.
11. Gray, M.W. et al. (2010) Cell biology. Irremediable complexity? *Science* 330 (6006), 920-921.
12. Ochsenreiter, T. et al. (2008) Alternative mRNA editing in trypanosomes is extensive and may contribute to mitochondrial protein diversity. *PLoS. ONE* 3 (2), e1566.

587 13. Ochsenreiter, T. et al. (2007) KISS: the kinetoplastid RNA editing sequence search tool.
588 RNA 13 (1), 1-4.

589 14. Kirby, L.E. et al. (2016) Analysis of the Trypanosoma brucei EATRO 164 Bloodstream
590 Guide RNA Transcriptome. PLoS Negl Trop Dis 10 (7), e0004793.

591 15. Koslowsky, D. et al. (2014) The insect-phase gRNA transcriptome in Trypanosoma
592 brucei. Nucleic Acids Res 42 (3), 1873-1886.

593 16. Aphasizhev, R. and Aphasizheva, I. (2014) Mitochondrial RNA editing in trypanosomes:
594 Small RNAs in control. Biochimie 100, 125-131.

595 17. Aphasizhev, R. and Aphasizheva, I. (2011) Uridine Insertion/Deletion mRNA Editing In
596 Trypanosomes: A Playground for RNA-guided Information Transfer. Wiley Interdisciplinary
597 Reviews: RNA 2, 669-685.

598 18. Hashimi, H. et al. (2013) Dual core processing: MRB1 is an emerging kinetoplast RNA
599 editing complex. Trends Parasitol 29 (2), 91-99.

600 19. Read, L.K. et al. (2016) Trypanosome RNA editing: the complexity of getting U in and
601 taking U out. Wiley Interdiscip Rev RNA 7 (1), 33-51.

602 20. Goring, H.U. (2012) 'Gestalt,' composition and function of the Trypanosoma brucei
603 editosome. Annu. Rev. Microbiol 66, 65-82.

604 21. Cruz-Reyes, J. et al. (2018) Dynamic RNA holo-editosomes with subcomplex variants:
605 Insights into the control of trypanosome editing. Wiley Interdiscip Rev RNA 9 (6), e1502.

606 22. Zimmer, S.L. et al. (2018) High throughput sequencing revolution reveals conserved
607 fundamentals of U-indel editing. Wiley Interdiscip Rev RNA, e1487.

608 23. Sement FM, S.T., Zhang L, Yu T, Huang L, Aphasizheva I, Aphasizhev R. (2018)
609 Transcription initiation defines kinetoplast RNA boundaries. PNAS 115 (44).

610 24. Suematsu, T. et al. (2016) Antisense Transcripts Delimit Exonucleolytic Activity of the
611 Mitochondrial 3' Processome to Generate Guide RNAs. Mol Cell 61 (3), 364-78.

612 25. Aphasizheva, I. and Aphasizhev, R. (2010) RET1-catalyzed Uridylylation Shapes the
613 Mitochondrial Transcriptome in *Trypanosoma brucei*. *Molecular and Cellular Biology* 30 (6),
614 1555-1567.

615 26. Mattiaccio, J.L. and Read, L.K. (2008) Roles for TbDSS-1 in RNA surveillance and decay
616 of maturation by-products from the 12S rRNA locus. *Nucleic Acids Res* 36 (1), 319-329.

617 27. Ryan, C.M. and Read, L.K. (2005) UTP-dependent turnover of *Trypanosoma brucei*
618 mitochondrial mRNA requires UTP polymerization and involves the RET1 TUTase. *RNA* 11
619 (5), 763-773.

620 28. Kao, C.Y. and Read, L.K. (2005) Opposing effects of polyadenylation on the stability of
621 edited and unedited mitochondrial RNAs in *Trypanosoma brucei*. *Mol. Cell Biol* 25 (5),
622 1634-1644.

623 29. Zhang, L. et al. (2017) PPR polyadenylation factor defines mitochondrial mRNA identity
624 and stability in trypanosomes. *EMBO J* 36, 2435-2454.

625 30. Aphasizheva, I. et al. (2011) Pentatricopeptide Repeat Proteins Stimulate mRNA
626 Adenylation/Uridylation to Activate Mitochondrial Translation in Trypanosomes. *Molecular*
627 *Cell* 42 (1), 106-117.

628 31. Etheridge, R.D. et al. (2008) 3' adenylation determines mRNA abundance and monitors
629 completion of RNA editing in *T. brucei* mitochondria. *EMBO J* 27 (11), 1596-1608.

630 32. Mesitov, M.V. et al. (2019) Pentatricopeptide repeat poly(A) binding protein KPAF4
631 stabilizes mitochondrial mRNAs in *Trypanosoma brucei*. *Nat Commun* 10 (1), 146.

632 33. Aphasizheva, I. et al. (2016) Ribosome-associated pentatricopeptide repeat proteins
633 function as translational activators in mitochondria of trypanosomes. *Mol Microbiol* 99 (6),
634 1043-58.

635 34. Ridlon, L. et al. (2013) The Importance of the 45S Ribosomal Small Subunit-related
636 Complex for Mitochondrial Translation in *Trypanosoma brucei*. J. Biol. Chem 288, 32963-
637 32978.

638 35. Saurer, M. et al. (2019) Mitoribosomal small subunit biogenesis in trypanosomes involves
639 an extensive assembly machinery. Science 365 (6458), 1144-1149.

640 36. Stuart, K.D. et al. (2005) Complex management: RNA editing in trypanosomes. Trends
641 Biochem. Sci 30 (2), 97-105.

642 37. Simpson, L. et al. (2010) Guide to the nomenclature of kinetoplastid RNA editing: a
643 proposal. Protist 161 (1), 2-6.

644 38. Ojala, D. et al. (1981) tRNA punctuation model of RNA processing in human
645 mitochondria. Nature 290, 470-474.

646 39. Read, L.K. et al. (1992) Extensive editing of both processed and preprocessed maxicircle
647 CR6 transcripts in *Trypanosoma brucei*. J. Biol. Chem 267, 1123-1128.

648 40. Koslowsky, D.J. and Yahampath, G. (1997) Mitochondrial mRNA 3' cleavage
649 polyadenylation and RNA editing in *Trypanosoma brucei* are independent events. Mol.
650 Biochem. Parasitol 90 (1), 81-94.

651 41. Michelotti, E.F. et al. (1992) *Trypanosoma brucei* mitochondrial ribosomal RNA
652 synthesis, processing and developmentally regulated expression. Mol. Biochem. Parasitol 54,
653 31-42.

654 42. Carnes, J. et al. (2015) Bloodstream form *Trypanosoma brucei* do not require mRPN1 for
655 gRNA processing. RNA 21 (1), 28-35.

656 43. Adler, B.K. et al. (1991) Modification of *Trypanosoma brucei* mitochondrial rRNA by
657 posttranscriptional 3' polyuridine tail formation. Mol. Cell. Biol 11, 5878-5884.

658 44. Bhat, G.J. et al. (1991) The two ATPase 6 mRNAs of *Leishmania tarentolae* differ at their
659 3' ends. Mol. Biochem. Parasitol 48, 139-150.

660 45. Blum, B. et al. (1990) A model for RNA editing in kinetoplastid mitochondria: "Guide"
 661 RNA molecules transcribed from maxicircle DNA provide the edited information. *Cell* 60,
 662 189-198.

663 46. Madina, B.R. et al. (2011) Guide RNA biogenesis involves a novel RNase III family
 664 endoribonuclease in *Trypanosoma brucei*. *RNA* 17 (10), 1821-30.

665 47. Schnauffer, A. et al. (2001) An RNA ligase essential for RNA editing and survival of the
 666 bloodstream form of *Trypanosoma brucei*. *Science* 291, 2159-2161.

667 48. Aphasizhev, R. et al. (2003) A tale of two TUTases. *Proc. Natl. Acad. Sci. U. S. A* 100
 668 (19), 10617-10622.

669 49. Aphasizhev, R. et al. (2002) Trypanosome Mitochondrial 3' Terminal Uridylyl
 670 Transferase (TUTase): The Key Enzyme in U-insertion/deletion RNA Editing. *Cell* 108, 637-
 671 648.

672 50. Mattiaccio, J.L. and Read, L.K. (2009) Evidence for a degradosome-like complex in the
 673 mitochondria of *Trypanosoma brucei*. *FEBS Lett* 583 (14), 2333-2338.

674 51. Aphasizheva, I. et al. (2004) RNA-editing terminal uridylyl transferase 1: identification of
 675 functional domains by mutational analysis. *J. Biol. Chem* 279 (23), 24123-24130.

676 52. Aphasizheva, I. et al. (2014) RNA binding and core complexes constitute the U-
 677 insertion/deletion editosome. *Mol. Cell Biol* 34 (23), 4329-4342.

678 53. Aphasizhev, R. et al. (2016) Constructive edge of uridylation-induced RNA degradation.
 679 *RNA Biol* 13 (11), 1078-1083.

680 54. Menezes, M.R. et al. (2018) 3' RNA Uridylation in Epitranscriptomics, Gene Regulation,
 681 and Disease. *Front Mol Biosci* 5, 61.

682 55. Jasmer, D. and Stuart, K. (1986) Sequence organization in African trypanosome
 683 minicircles is defined by 18 base pair inverted repeats. *Mol. Biochem. Parasitol* 18, 321-332.

684 56. Pollard, V.W. et al. (1990) Organization of minicircle genes for guide RNAs in
685 *Trypanosoma brucei*. Cell 63, 783-790.

686 57. Cooper, S. et al. (2019) Assembly and annotation of the mitochondrial minicircle genome
687 of a differentiation-competent strain of *Trypanosoma brucei*. Nucleic Acids Res.

688 58. Simpson, L. et al. (2015) Comparison of the Mitochondrial Genomes and Steady State
689 Transcriptomes of Two Strains of the Trypanosomatid Parasite, *Leishmania tarentolae*. Plos
690 Neglected Tropical Diseases 9 (7).

691 59. Simpson, R.M. et al. (2016) High-throughput sequencing of partially edited trypanosome
692 mRNAs reveals barriers to editing progression and evidence for alternative editing. RNA 22
693 (5), 677-95.

694 60. Chung, C.Z. et al. (2017) Tipping the balance of RNA stability by 3' editing of the
695 transcriptome. Biochim Biophys Acta Gen Subj 1861 (11 Pt B), 2971-2979.

696 61. Leung, S.S. and Koslowsky, D.J. (2001) RNA editing in *Trypanosoma brucei*:
697 characterization of gRNA U-tail interactions with partially edited mRNA substrates. Nucleic
698 Acids Res 29 (3), 703-709.

699 62. Arts, G.J. et al. (1995) A possible role for the guide RNA U-tail as a specificity
700 determinant in formation of guide RNA-messenger RNA chimeras in mitochondrial extracts
701 of *Crithidia fasciculata*. Mol. Biochem. Parasitol 73 (1-2), 211-222.

702 63. Blum, B. and Simpson, L. (1990) Guide RNAs in kinetoplastid mitochondria have a
703 nonencoded 3' oligo-(U) tail involved in recognition of the pre-edited region. Cell 62, 391-
704 397.

705 64. Ryan, C.M. et al. (2003) Polyadenylation regulates the stability of *Trypanosoma brucei*
706 mitochondrial RNAs. J. Biol. Chem 278 (35), 32753-32762.

707 65. Small, I.D. and Peeters, N. (2000) The PPR motif - a TPR-related motif prevalent in plant
708 organellar proteins. Trends Biochem Sci 25 (2), 46-47.

709 66. Shen, C. et al. (2016) Structural basis for specific single-stranded RNA recognition by
710 designer pentatricopeptide repeat proteins. *Nat Commun* 7, 11285.

711 67. Barkan, A. and Small, I. (2014) Pentatricopeptide repeat proteins in plants. *Annu Rev*
712 *Plant Biol* 65, 415-42.

713 68. Pfalz, J. et al. (2009) Site-specific binding of a PPR protein defines and stabilizes 5' and 3'
714 mRNA termini in chloroplasts. *EMBO J* 28 (14), 2042-2052.

715 69. Cheng, S. et al. (2016) Redefining the structural motifs that determine RNA binding and
716 RNA editing by pentatricopeptide repeat proteins in land plants. *Plant J* 85 (4), 532-47.

717 70. Feagin, J.E. et al. (1988) Creation of AUG initiation codons by addition of uridines within
718 cytochrome b transcripts of kinetoplastids. *PNAS* 85, 539-543.

719 71. Koslowsky, D.J. et al. (1990) The MURF3 gene of *T. brucei* contains multiple domains of
720 extensive editing and is homologous to a subunit of NADH dehydrogenase. *Cell* 62, 901-911.

721 72. Seiwert, S.D. et al. (1996) Direct visualization of uridylyte deletion in vitro suggests a
722 mechanism for kinetoplastid RNA editing. *Cell* 84, 831-841.

723 73. Kable, M.L. et al. (1996) RNA editing: a mechanism for gRNA-specified uridylyte
724 insertion into precursor mRNA [see comments]. *Science* 273, 1189-1195.

725 74. Seiwert, S.D. and Stuart, K. (1994) RNA editing: transfer of genetic information from
726 gRNA to precursor mRNA in vitro. *Science* 266, 114-117.

727 75. Sturm, N.R. and Simpson, L. (1990) Kinetoplast DNA minicircles encode guide RNAs
728 for editing of cytochrome oxidase subunit III mRNA. *Cell* 61, 879-884.

729 76. Golden, D.E. and Hajduk, S.L. (2005) The 3'-untranslated region of cytochrome oxidase
730 II mRNA functions in RNA editing of African trypanosomes exclusively as a cis guide RNA.
731 *RNA* 11 (1), 29-37.

732 77. Koslowsky, D.J. et al. (1991) Cycles of progressive realignment of gRNA with mRNA in
733 RNA editing. *Cell* 67, 537-546.

734 78. Simpson, R.M. et al. (2017) Trypanosome RNA Editing Mediator Complex proteins have
735 distinct functions in gRNA utilization. *Nucleic Acids Res* 45 (13), 7965-7983.

736 79. Gerasimov, E.S. et al. (2018) Trypanosomatid mitochondrial RNA editing: dramatically
737 complex transcript repertoires revealed with a dedicated mapping tool. *Nucleic Acids Res* 46
738 (2), 765-781.

739 80. Maslov, D.A. and Simpson, L. (1992) The polarity of editing within a multiple gRNA-
740 mediated domain is due to formation of anchors for upstream gRNAs by downstream editing.
741 *Cell* 70, 459-467.

742 81. Igo, R.P., Jr. et al. (2002) RNA sequence and base pairing effects on insertion editing in
743 *Trypanosoma brucei*. *Mol Cell Biol* 22 (5), 1567-1576.

744 82. Blanc, V. et al. (1999) The mitochondrial RNA ligase from *Leishmania tarentolae* can
745 join RNA molecules bridged by a complementary RNA. *J. Biol. Chem* 274, 24289-24296.

746 83. Cruz-Reyes, J. et al. (2001) Trypanosome RNA editing: simple guide RNA features
747 enhance U deletion 100-fold. *Mol. Cell Biol* 21 (3), 884-892.

748 84. Leung, S.S. and Koslowsky, D.J. (2001) Interactions of mRNAs and gRNAs involved in
749 trypanosome mitochondrial RNA editing: Structure probing of an mRNA bound to its
750 cognate gRNA. *RNA* 7 (12), 1803-1816.

751 85. Aphasizhev, R. et al. (2003) Isolation of a U-insertion/deletion editing complex from
752 *Leishmania tarentolae* mitochondria. *EMBO J* 22 (4), 913-924.

753 86. Panigrahi, A.K. et al. (2003) Identification of novel components of *Trypanosoma brucei*
754 editosomes. *RNA* 9 (4), 484-492.

755 87. Golas, M.M. et al. (2009) Snapshots of the RNA editing machine in trypanosomes
756 captured at different assembly stages in vivo. *EMBO J* 28 (6), 766-778.

757 88. Li, F. et al. (2009) Structure of the core editing complex (L-complex) involved in uridine
758 insertion/deletion RNA editing in trypanosomatid mitochondria. *Proc. Natl. Acad. Sci. U. S.*
759 *A* 106 (30), 12306-12310.

760 89. Rusche, L.N. et al. (1997) Purification of a functional enzymatic editing complex from
761 *Trypanosoma brucei* mitochondria. *EMBO J* 16 (13), 4069-4081.

762 90. Schnauffer, A. et al. (2003) Separate Insertion and Deletion Subcomplexes of the
763 *Trypanosoma brucei* RNA Editing Complex. *Mol Cell* 12, 307-319.

764 91. Tarun, S.Z., Jr. et al. (2008) KREPA6 is an RNA-binding protein essential for editosome
765 integrity and survival of *Trypanosoma brucei*. *RNA* 14 (2), 347-358.

766 92. Guo, X. et al. (2008) The KREPA3 zinc finger motifs and OB-fold domain are essential
767 for RNA editing and survival of *Trypanosoma brucei*. *Mol Cell Biol* 28 (22), 6939-6953.

768 93. Babbarwal, V.K. et al. (2007) An essential role of KREPB4 in RNA editing and structural
769 integrity of the editosome in *Trypanosoma brucei*. *RNA* 13 (5), 737-744.

770 94. Salavati, R. et al. (2006) KREPA4, an RNA binding protein essential for editosome
771 integrity and survival of *Trypanosoma brucei*. *RNA* 12 (5), 819-831.

772 95. Guo, X. et al. (2012) KREPB6, KREPB7, and KREPB8 are important for editing
773 endonuclease function in *Trypanosoma brucei*. *RNA* 18 (2), 308-320.

774 96. Carnes, J. et al. (2012) KREX2 is not essential for either procyclic or bloodstream form
775 *Trypanosoma brucei*. *PLoS. ONE* 7 (3), e33405.

776 97. Ernst, N.L. et al. (2009) Differential functions of two editosome exoUases in
777 *Trypanosoma brucei*. *RNA* 15 (5), 947-957.

778 98. Carnes, J. et al. (2008) RNA Editing in *Trypanosoma brucei* requires three different
779 editosomes. *Mol. Cell Biol* 28 (1), 122-130.

780 99. Carnes, J. et al. (2017) In vivo cleavage specificity of *Trypanosoma brucei* editosome
781 endonucleases. *Nucleic Acids Res* 45 (8), 4667-4686.

782 100. Macrae, I.J. and Doudna, J.A. (2007) Ribonuclease revisited: structural insights into
783 ribonuclease III family enzymes. *Curr. Opin. Struct. Biol* 17 (1), 138-145.

784 101. Carnes, J. et al. (2012) Mutational analysis of *Trypanosoma brucei* editosome proteins
785 KREPB4 and KREPB5 reveals domains critical for function. *RNA* 18 (10), 1897-1909.

786 102. McDermott, S.M. et al. (2019) Editosome RNase III domain interactions are essential
787 for editing and differ between life cycle stages in *Trypanosoma brucei*. *RNA* 25 (9), 1150-
788 1163.

789 103. McDermott, S.M. and Stuart, K. (2017) The essential functions of KREPB4 are
790 developmentally distinct and required for endonuclease association with editosomes. *RNA* 23
791 (11), 1672-1684.

792 104. Carnes, J. et al. (2011) Endonuclease associations with three distinct editosomes in
793 *Trypanosoma brucei*. *J. Biol. Chem* 286 (22), 19320-19330.

794 105. Panigrahi, A.K. et al. (2006) Compositionally and functionally distinct editosomes in
795 *Trypanosoma brucei*. *RNA* 12 (6), 1038-1049.

796 106. McDermott, S.M. et al. (2016) The Architecture of *Trypanosoma brucei* editosomes.
797 *Proc Natl Acad Sci U S A* 113 (42), E6476-E6485.

798 107. Schnauffer, A. et al. (2010) A protein-protein interaction map of trypanosome ~20S
799 editosomes. *J. Biol. Chem* 285 (8), 5282-5295.

800 108. Kang, X. et al. (2006) Reconstitution of full-round uridine-deletion RNA editing with
801 three recombinant proteins. *Proc. Natl. Acad. Sci. U. S. A* 103 (38), 13944-13949.

802 109. Gao, G. et al. (2005) Functional complementation of *Trypanosoma brucei* RNA in vitro
803 editing with recombinant RNA ligase. *Proc. Natl. Acad. Sci. U. S. A* 102 (13), 4712-4717.

804 110. Rogers, K. et al. (2007) Uridylate-specific 3' 5'-exoribonucleases involved in uridylate-
805 deletion RNA editing in trypanosomatid mitochondria. *J. Biol. Chem* 282 (40), 29073-29080.

806 111. Ringpis, G.E. et al. (2010) Mechanism of U Insertion RNA Editing in Trypanosome
807 Mitochondria: The Bimodal TUTase Activity of the Core Complex. Journal of Molecular
808 Biology 399 (5), 680-695.

809 112. Ringpis, G.E. et al. (2010) Mechanism of U-insertion RNA Editing in Trypanosome
810 Mitochondria: Characterization of RET2 Functional Domains by Mutational Analysis. J. Mol
811 Biol 399 (5), 696-706.

812 113. Ernst, N.L. et al. (2003) TbMP57 is a 3' terminal uridylyl transferase (TUTase) of the
813 Trypanosoma brucei editosome. Mol. Cell 11 (6), 1525-1536.

814 114. Deng, J. et al. (2005) Structural basis for UTP specificity of RNA editing TUTases from
815 Trypanosoma brucei. EMBO J 24 (23), 4007-4017.

816 115. Gao, G. and Simpson, L. (2003) Is the Trypanosoma brucei REL1 RNA ligase specific
817 for U-deletion RNA editing, and is the REL2 RNA ligase specific for U-insertion editing? J.
818 Biol. Chem 278 (30), 27570-27574.

819 116. Huang, C.E. et al. (2001) Roles for ligases in the RNA editing complex of *Trypanosoma*
820 *brucei*: band IV is needed for U-deletion and RNA repair. EMBO Journal 20 (17), 4694-
821 4703.

822 117. Pollard, V.W. et al. (1992) Native mRNA editing complexes from Trypanosoma brucei
823 mitochondria. EMBO J 11, 4429-4438.

824 118. Osato, D. et al. (2009) Uridine insertion/deletion RNA editing in trypanosomatid
825 mitochondria: In search of the editosome. RNA 15 (7), 1338-1344.

826 119. Aphasizheva, I. and Aphasizhev, R. (2015) U-Insertion/Deletion mRNA-Editing
827 Holoenzyme: Definition in Sight. Trends Parasitol 13 (11), 1078-1083.

828 120. Panigrahi, A.K. et al. (2007) Mitochondrial complexes in trypanosoma brucei: a novel
829 complex and a unique oxidoreductase complex. Mol. Cell Proteomics 7 (3), 534-545.

830 121. Weng, J. et al. (2008) Guide RNA-Binding Complex from Mitochondria of
831 Trypanosomatids. *Molecular Cell* 32, 198-209.

832 122. Hashimi, H. et al. (2009) Kinetoplastid guide RNA biogenesis is dependent on subunits
833 of the mitochondrial RNA binding complex 1 and mitochondrial RNA polymerase. *RNA* 15
834 (4), 588-599.

835 123. Madina, B.R. et al. (2014) Native mitochondrial RNA-binding complexes in
836 kinetoplastid RNA editing differ in guide RNA composition. *RNA* 20 (7), 1142-1152.

837 124. Huang, Z. et al. (2015) Integrity of the core mitochondrial RNA-binding complex 1 is
838 vital for trypanosome RNA editing. *RNA* 21 (12), 2088-102.

839 125. McAdams, N.M. et al. (2019) MRB10130 is a RESC assembly factor that promotes
840 kinetoplastid RNA editing initiation and progression. *RNA* 25 (9), 1177-1191.

841 126. McAdams, N.M. et al. (2018) MRB7260 is essential for productive protein-RNA
842 interactions within the RNA editing substrate binding complex during trypanosome RNA
843 editing. *RNA* 24 (4), 540-556.

844 127. Ammerman, M.L. et al. (2010) TbRGG2 facilitates kinetoplastid RNA editing initiation
845 and progression past intrinsic pause sites. *RNA* 16 (11), 2239-2251.

846 128. Dixit, S. et al. (2017) Differential Binding of Mitochondrial Transcripts by MRB8170
847 and MRB4160 Regulates Distinct Editing Fates of Mitochondrial mRNA in Trypanosomes.
848 *MBio* 8 (1).

849 129. Ammerman, M.L. et al. (2012) Architecture of the trypanosome RNA editing accessory
850 complex, MRB1. *Nucleic Acids Res* 40 (12), 5637-5650.

851 130. Aphasizhev, R. et al. (2003) A 100-kD complex of two RNA-binding proteins from
852 mitochondria of *Leishmania tarentolae* catalyzes RNA annealing and interacts with several
853 RNA editing components. *RNA* 9 (1), 62-76.

854 131. Nikpour, N. and Salavati, R. (2019) The RNA binding activity of the first identified
855 trypanosome protein with Z-DNA-binding domains. *Sci Rep* 9 (1), 5904.

856 132. Kumar, V. et al. (2019) Protein features for assembly of the RNA editing helicase 2
857 subcomplex (REH2C) in Trypanosome holo-editosomes. *PLoS One* 14 (4), e0211525.

858 133. Madina, B.R. et al. (2015) Native Variants of the MRB1 Complex Exhibit Specialized
859 Functions in Kinetoplastid RNA Editing. *PLoS. ONE* 10 (4), e0123441.

860 134. Hernandez, A. et al. (2010) REH2 RNA helicase in kinetoplastid mitochondria:
861 ribonucleoprotein complexes and essential motifs for unwinding and guide RNA (gRNA)
862 binding. *J. Biol. Chem* 285 (2), 1220-1228.

863 135. Kumar, V. et al. (2016) REH2C Helicase and GRBC Subcomplexes May Base Pair
864 through mRNA and Small Guide RNA in Kinetoplastid Editosomes. *J Biol Chem* 291 (11),
865 5753-64.

866 136. Kao, C.Y. and Read, L.K. (2007) Targeted depletion of a mitochondrial
867 nucleotidyltransferase suggests the presence of multiple enzymes that polymerize mRNA 3'
868 tails in *Trypanosoma brucei* mitochondria. *Mol. Biochem. Parasitol* 154 (2), 158-169.

869 137. Missel, A. et al. (1995) A putative RNA helicase of the DEAD box family from
870 *Trypanosoma brucei*. *Mol. Biochem. Parasitol* 75 (1), 123-126.

871 138. Missel, A. et al. (1997) Disruption of a gene encoding a novel mitochondrial DEAD-box
872 protein in *Trypanosoma brucei* affects edited mRNAs. *Molecular and Cellular Biology* 17
873 (9), 4895-4903.

874 139. Li, F. et al. (2011) Trypanosome REH1 is an RNA helicase involved with the 3'-5'
875 polarity of multiple gRNA-guided uridine insertion/deletion RNA editing. *Proc. Natl. Acad.*
876 *Sci. U. S. A* 108 (9), 3542-3547.

877 140. Koller, J. et al. (1997) *Trypanosoma brucei* gBP21. An arginine-rich mitochondrial
878 protein that binds to guide RNA with high affinity. *J. Biol. Chem* 272, 3749-3757.

879 141. Blom, D. et al. (2001) Cloning and characterization of two guide RNA-binding proteins
880 from mitochondria of *Crithidia fasciculata*: gBP27, a novel protein, and gBP29, the
881 orthologue of *Trypanosoma brucei* gBP21. *Nucleic Acids Res* 29 (14), 2950-2962.

882 142. Zikova, A. et al. (2008) Structure and function of the native and recombinant
883 mitochondrial MRP1/MRP2 complex from *Trypanosoma brucei*. *Int. J. Parasitol.*

884 143. Schumacher, M.A. et al. (2006) Crystal structures of *T. brucei* MRP1/MRP2 guide-RNA
885 binding complex reveal RNA matchmaking mechanism. *Cell* 126 (4), 701-711.

886 144. Muller, U.F. and Goring, H.U. (2002) Mechanism of the gBP21-mediated RNA/RNA
887 annealing reaction: matchmaking and charge reduction. *Nucleic Acids Res* 30 (2), 447-455.

888 145. Muller, U.F. et al. (2001) Annealing of RNA editing substrates facilitated by guide
889 RNA-binding protein gBP21. *EMBO J* 20 (6), 1394-1404.

890 146. Fisk, J.C. et al. (2009) Distinct and overlapping functions of MRP1/2 and RBP16 in
891 mitochondrial RNA metabolism. *Mol. Cell Biol* 29 (19), 5214-5225.

892 147. Vondruskova, E. et al. (2005) RNA interference analyses suggest a transcript-specific
893 regulatory role for mitochondrial RNA-binding proteins MRP1 and MRP2 in RNA editing
894 and other RNA processing in *Trypanosoma brucei*. *J. Biol. Chem* 280 (4), 2429-2438.

895 148. Tylec, B.L. et al. (2019) Intrinsic and regulated properties of minimally edited
896 trypanosome mRNAs. *Nucleic Acids Res* 47 (7), 3640-3657.

897 149. Hayman, M.L. and Read, L.K. (1999) *Trypanosoma brucei* RBP16 is a mitochondrial Y-
898 box family protein with guide RNA binding activity. *J. Biol. Chem* 274 (17), 12067-12074.

899 150. Ammerman, M.L. et al. (2008) gRNA/pre-mRNA annealing and RNA chaperone
900 activities of RBP16. *RNA*.

901 151. Miller, M.M. et al. (2006) RBP16 stimulates trypanosome RNA editing in vitro at an
902 early step in the editing reaction. *RNA* 12 (7), 1292-1303.

903 152. Pelletier, M. and Read, L.K. (2003) RBP16 is a multifunctional gene regulatory protein
 904 involved in editing and stabilization of specific mitochondrial mRNAs in *Trypanosoma*
 905 *brucei*. RNA 9 (4), 457-468.

906 153. Miller, M.M. and Read, L.K. (2003) *Trypanosoma brucei*: functions of RBP16 cold
 907 shock and RGG domains in macromolecular interactions. Exp. Parasitol 105 (2), 140-148.

908 154. Hayman, M.L. et al. (2001) The trypanosome homolog of human p32 interacts with
 909 RBP16 and stimulates its gRNA binding activity. Nucleic Acids Res 29 (24), 5216-5225.

910 155. Pelletier, M. et al. (2000) RNA-binding properties of the mitochondrial Y-box protein
 911 RBP16. Nucleic Acids Res 28 (5), 1266-1275.

912 156. Vanhamme, L. et al. (1998) *Trypanosoma brucei* TBRGG1, a mitochondrial oligo(U)-
 913 binding protein that co-localizes with an *in vitro* RNA editing activity. Journal of Biological
 914 Chemistry 273 (34), 21825-21833.

915 157. Hashimi, H. et al. (2008) TbRGG1, an essential protein involved in kinetoplastid RNA
 916 metabolism that is associated with a novel multiprotein complex. RNA 14, 970-980.

917 158. McAdams, N.M. et al. (2015) An arginine-glycine-rich RNA binding protein impacts
 918 the abundance of specific mRNAs in the mitochondria of *Trypanosoma brucei*. Eukaryot Cell
 919 14 (2), 149-57.

920 159. Shaw, P.L. et al. (2015) Structures of the *T. brucei* kRNA editing factor MRB1590
 921 reveal unique RNA-binding pore motif contained within an ABC-ATPase fold. Nucleic Acids
 922 Res 43 (14), 7096-109.

923 160. Madison-Antenucci, S. and Hajduk, S.L. (2001) RNA editing-associated protein 1 is an
 924 RNA binding protein with specificity for preedited mRNA. Mol Cell 7 (4), 879-886.

925 161. Madison-Antenucci, S. et al. (1998) Kinetoplastid RNA-editing-associated protein 1
 926 (REAP-1): a novel editing complex protein with repetitive domains. EMBO J 17, 6368-6376.

927 162. Hans, J. et al. (2007) RNA-editing-associated protein 1 null mutant reveals link to
 928 mitochondrial RNA stability. *RNA* 13 (6), 881-889.

929 163. Dixit, S. and Lukes, J. (2018) Combinatorial interplay of RNA-binding proteins tunes
 930 levels of mitochondrial mRNA in trypanosomes. *RNA* 24 (11), 1594-1606.

931 164. Zimmer, S.L. et al. (2011) A novel member of the RNase D exoribonuclease family
 932 functions in mitochondrial guide RNA metabolism in *Trypanosoma brucei*. *J. Biol. Chem*
 933 286 (12), 10329-10340.

934 165. Taschner, A. et al. (2012) Nuclear RNase P of *Trypanosoma brucei*: a single protein in
 935 place of the multicomponent RNA-protein complex. *Cell Rep* 2 (1), 19-25.

936 166. Salavati, R. et al. (2001) Mitochondrial ribonuclease P activity of *Trypanosoma brucei*.
 937 *Molecular and Biochemical Parasitology* 115 (1), 109-117.

938 167. Kapushoc, S.T. et al. (2002) Differential localization of nuclear-encoded tRNAs
 939 between the cytosol and mitochondrion in *Leishmania tarentolae*. *RNA* 8 (1), 57-68.

940 168. Kapushoc, S.T. et al. (2000) End processing precedes mitochondrial importation and
 941 editing of tRNAs in *Leishmania tarentolae*. *J. Biol Chem* 275 (48), 37907-37914.

942 169. Rajappa-Titu, L. et al. (2016) RNA Editing TUTase 1: structural foundation of substrate
 943 recognition, complex interactions and drug targeting. *Nucleic Acids Res* 44 (22), 10862-
 944 10878.

945 170. Mingler, M.K. et al. (2006) Identification of pentatricopeptide repeat proteins in
 946 *Trypanosoma brucei*. *Mol. Biochem. Parasitol* 150 (1), 37-45.

947 171. Pusnik, M. et al. (2007) Pentatricopeptide repeat proteins in *Trypanosoma brucei*
 948 function in mitochondrial ribosomes. *Mol. Cell Biol* 27 (19), 6876-6888.

949 172. Trotter, J.R. et al. (2005) A deletion site editing endonuclease in *Trypanosoma brucei*.
 950 *Mol. Cell* 20 (3), 403-412.

951 173. Carnes, J. et al. (2005) An essential RNase III insertion editing endonuclease in
 952 *Trypanosoma brucei*. *Proc. Natl. Acad. Sci. U. S. A* 102 (46), 16614-16619.
 953 174. Drozdz, M. et al. (2002) TbMP81 is required for RNA editing in *Trypanosoma brucei*.
 954 *EMBO J* 21 (7), 1791-1799.
 955 175. Panigrahi, A.K. et al. (2001) Four Related Proteins of the *Trypanosoma brucei* RNA
 956 Editing Complex. *Mol Cell Biol* 21 (20), 6833-6840.
 957 176. McDermott, S.M. et al. (2015) Differential Editosome Protein Function between Life
 958 Cycle Stages of *Trypanosoma brucei*. *J. Biol. Chem.*
 959 177. Law, J.A. et al. (2007) In *Trypanosoma brucei* RNA editing, TbMP18 (band VII) is
 960 critical for editosome integrity and for both insertional and deletional cleavages. *Mol. Cell*
 961 *Biol* 27 (2), 777-787.
 962 178. Wang, B. et al. (2003) TbMP44 is essential for RNA editing and structural integrity of
 963 the editosome in *Trypanosoma brucei*. *Eukaryot Cell* 2 (3), 578-87.
 964 179. Lerch, M. et al. (2012) Editosome accessory factors KREPB9 and KREPB10 in
 965 *Trypanosoma brucei*. *Eukaryot. Cell* 11 (7), 832-843.
 966 180. Carnes, J. et al. (2018) RNase III Domain of KREPB9 and KREPB10 Association with
 967 Editosomes in *Trypanosoma brucei*. *mSphere* 3 (1).
 968 181. Aphasizheva, I. et al. (2009) Novel TUTase associates with an editosome-like complex
 969 in mitochondria of *Trypanosoma brucei*. *RNA* 15, 1322-1337.
 970 182. Acestor, N. et al. (2009) The MRB1 complex functions in kinetoplastid RNA
 971 processing. *RNA* 15 (2), 277-286.
 972 183. Ammerman, M.L. et al. (2013) A core MRB1 complex component is indispensable for
 973 RNA editing in insect and human infective stages of *Trypanosoma brucei*. *PLoS. ONE* 8
 974 (10), e78015.

975 184. Ammerman, M.L. et al. (2011) MRB3010 is a core component of the MRB1 complex
 976 that facilitates an early step of the kinetoplastid RNA editing process. *RNA* 17 (5), 865-877.
 977 185. Kafkova, L. et al. (2012) Functional characterization of two paralogs that are novel RNA
 978 binding proteins influencing mitochondrial transcripts of *Trypanosoma brucei*. *RNA* 18 (10),
 979 1846-1861.
 980 186. Fisk, J.C. et al. (2008) TbRGG2, an essential RNA editing accessory factor in two
 981 *Trypanosoma brucei* life cycle stages. *J. Biol. Chem* 283 (34), 23016-23025.
 982 187. Foda, B.M. et al. (2012) Multifunctional G-rich and RRM-containing domains of
 983 TbRGG2 perform separate yet essential functions in trypanosome RNA editing. *Eukaryot*
 984 *Cell* 11 (9), 1119-31.
 985 188. Travis, B. et al. (2019) The RRM of the kRNA-editing protein TbRGG2 uses multiple
 986 surfaces to bind and remodel RNA. *Nucleic Acids Res* 47 (4), 2130-2142.
 987 189. Missel, A. et al. (1997) Disruption of a gene encoding a novel mitochondrial DEAD-box
 988 protein in *Trypanosoma brucei* affects edited mRNAs. *Mol Cell Biol* 17 (9), 4895-4903.
 989
 990

991 **Glossary**

992 **5' and 3' cleavage fragments:** mRNA fragments generated by gRNA-directed
993 endonucleolytic cleavage.

994 **Anchor:** 5' part of the gRNA that forms a continuous 10-15 nt duplex with pre-edited,
995 partially-edited or fully-edited mRNA; this region is responsible for initial gRNA-mRNA
996 interaction.

997 **Cryptogene:** a maxicircle gene with defective coding sequence; the defects are corrected by
998 U-insertion/deletion editing with concomitant restoration of protein reading frame.

999 **Editing block:** an mRNA segment covered by a single gRNA; often contains both U-
1000 insertion and U-deletion sites.

1001 **Editing domain:** an mRNA region covered by single or multiple overlapping gRNAs. In a
1002 multi-gRNA domain, sequence changes directed by the initiating gRNA create the binding
1003 site for the subsequent one. The hierarchical gRNA binding provides for the overall 3'-5'
1004 progression of editing events within a domain.

1005 **Editing site:** position of the gRNA-directed internal cleavage where uridines are either
1006 deleted from or inserted into the mRNA.

1007 **EEP domain:** endonuclease/exonuclease/phosphatase (EEP) domain in U-specific editing
1008 exonucleases.

1009 **Fully-edited mRNA:** a final editing product; contains a protein coding frame.

1010 **Guide RNA (gRNA):** a small non-coding RNA that specifies positions and extent of U-
1011 insertions and deletions by forming an imperfect duplex with pre-edited or partially-edited
1012 mRNA. Guide RNA is typically 30-60 nucleotides (nt) in length and possesses a 5'
1013 triphosphate and a 1-20 nt 3' U-tail.

1014 **Junction:** A region present in most partially-edited mRNAs at the 5' leading edge of editing;
1015 often displays mis-edited and non-canonically edited sequences. Junction-containing
1016 transcripts may represent intermediates that will be re-edited to canonical sequence, dead-end
1017 by-products, and mRNAs with a non-canonical protein coding sequence.

1018 **Kinetoplastids (class Kinetoplastea):** flagellated protists characterized by the presence of a
1019 kinetoplast. Phylogenetically positioned within the phylum Euglenozoa, this group includes
1020 the obligatory parasitic trypanosomatids (family *Trypanosomatidae*), free-living and parasitic
1021 bodonids, and more distantly related taxa.

1022 **Kinetoplast:** a densely packed nucleoprotein structure, disc-shaped and catenated in
1023 trypanosomatids, and dispersed to various degrees in most bodonids, that encloses
1024 mitochondrial DNA (kinetoplast DNA; kDNA). A non-dividing *T. brucei* cell contains a
1025 single mitochondrion with a single kinetoplast composed of catenated maxicircles (~23 kb
1026 each, few dozen copies) and minicircles (~1 kb each, ~5,000 units).

1027 **KPAC:** kinetoplast polyadenylation complex. A ribonucleoprotein complex of KPAP1
1028 poly(A) polymerase, and pentatricopeptide repeat-containing (PPR) RNA binding proteins
1029 designated Kinetoplast Polyadenylation Factors 1, 2, 3 and 4 (KPAF1, 2, 3 and 4).

1030 **Maxicircle:** an equivalent of a typical mitochondrial genome; includes a conserved ~15 kb
1031 region encoding 9S and 12S rRNAs, two guide (g)RNAs, and 18 protein genes. A variable
1032 region composed of repeated DNA sequences constitutes the rest of the molecule.

1033 **Minicircle:** the molecules forming the bulk of kinetoplast. Approximately 400 sequence
1034 classes present at various frequencies encode ~930 gRNAs required for the editing process
1035 and 370 gRNA-like molecules that likely participate in gRNA processing.

1036 **Moderately-edited mRNA:** a transcript with a few editing sites confined to a limited mRNA
1037 region.

1038 **MPsome:** mitochondrial 3' processome. A protein complex composed of Kinetoplast RNA
 1039 editing TUTase 1 (KRET1), 3'-5' exonuclease KDSS1, and MPSS1-6 subunits lacking
 1040 recognizable motifs.

1041 **MTRNAP:** mitochondrial RNA polymerase, a single-subunit T3/T7-like DNA-dependent
 1042 RNA polymerase.

1043 **Never-edited mRNA:** a maxicircle transcript containing an encoded open reading frame
 1044 which does not require editing.

1045 **Pan-edited mRNA:** a transcript that undergoes massive editing directed by multiple gRNAs.
 1046 There can be two editing domains within a pan-edited mRNA.

1047 **Partially-edited mRNA:** an intermediate of the editing process. Partially-edited mRNAs
 1048 often contain junctions whose sequences match neither pre-edited nor canonical fully-edited
 1049 mRNAs.

1050 **PPsome:** 5' pyrophosphate processome. Protein complex containing MERS1 NUDIX
 1051 pyrophosphohydrolase and MERS2 PPR RNA binding protein.

1052 **PPR:** Pentatricopeptide (35 amino acids) helix-turn-helix repeat. PPR arrays are present in
 1053 many trypanosomal mitochondrial RNA binding proteins.

1054 **Pre-edited mRNA:** a 3' processed monocistronic cryptogene transcript that must undergo
 1055 editing to acquire an open reading frame and/or translation initiation and termination signals.

1056 **RECC:** RNA Editing Catalytic Complex, formerly called ~20S editosome or RNA Editing
 1057 Core Complex. A protein complex of 14 or more subunits, depending on the isoform;
 1058 includes pre-mRNA cleavage, U-insertion, U-deletion and RNA ligation enzymes, and
 1059 structural and RNA binding factors.

1060 **RESC:** RNA Editing Substrate Binding Complex, formerly called Mitochondrial RNA
 1061 Binding Complex 1 (MRB1) and Guide RNA Binding Complex (GRBC). An ~20-subunit
 1062 modular protein complex that likely exists in several isoforms; most components lack
 1063 recognizable motifs. RESC binds RNA editing substrates, intermediates and products, and
 1064 coordinates interactions of gRNA and mRNA with RECC, REH2C and other auxiliary
 1065 factors during editing. RESC has also been implicated in coordination of pre-mRNA 5' and
 1066 3' modification processes.

1067 **REH2C:** RNA Editing Helicase 2 Complex, a protein complex formed by an ATP-dependent
 1068 DEAH/RHA RNA helicase KREH2, zinc finger protein KH2F1, and KH2F2 factor which
 1069 lacks recognizable motifs.

1070 **RNA editing holoenzyme (editosome):** a ribonucleoprotein particle consisting of RECC,
 1071 RESC and REH2C complexes, and several auxiliary factors.

1072 **RNA helicase:** a motor protein capable of harnessing the energy from NTP hydrolysis to
 1073 unwind double stranded RNAs or to remodel ribonucleoprotein complexes.

1074 **RNase II:** Exoribonuclease II cleaves single-stranded RNA in 3' to 5'-direction yielding
 1075 nucleoside 5' monophosphates.

1076 **RNase III:** Endoribonuclease III typically cleaves both strands in double-stranded RNA
 1077 leaving 5' monophosphate and 3' hydroxyl groups. RNase III editing endonucleases cleave
 1078 only the mRNA strand at an unpaired nucleotide adjacent to a gRNA-mRNA duplex.

1079 **TUTase:** terminal uridyltransferase, UTP-specific nucleotidyl transferase which adds U-
 1080 residues to the 3' end of RNA.

1081 **U-insertion/deletion mRNA editing:** a process by which U-residues are inserted into, or
1082 deleted from, a cryptogene transcript. Editing is directed by gRNAs and catalyzed by the
1083 RNA editing holoenzyme (editosome).

1084 **UTR:** untranslated region of mRNA.

1085

Table 1. Proposed nomenclature of mitochondrial RNA processing complexes and factors.

Gene identification numbers refer to *T. brucei* strain TREU927 predicted protein sequences

(TriTrypDB, Release 45, September 5, 2019, <https://tritrypdb.org/tritrypdb/>).

Legacy	Assigned	Function	Motifs	TriTryp ID	References
Nucleolytic Processing: Mitochondrial 3' processome (MPsome)					
RET1	KRET1	KRET1	TUTase, 3' uridylation of primary and mature RNAs	TUTase, PAP associated	Tb927.7.3950 [24, 25, 48, 49, 169]
KDSS1	KDSS1	KDSS1	3'-5' exonuclease	RNB (Ribonuclease II)	Tb927.9.7210 [24, 26, 50]
MPSS1		MPSS1			Tb927.11.9150 [24]
MPSS2		MPSS2			Tb927.10.9000 [24]
MPSS3		MPSS3			Tb927.3.2770 [24]
MPSS4		MPSS4			Tb927.10.6170 [24]
MPSS5		MPSS5			Tb927.9.4810 [24]
MPSS6		MPSS6			Tb927.6.2190 [24]
Modification of the 5' end: Pyrophosphohydrolase complex (PPsome)					
MERS1		MERS1	PPi removal from 5' end	NUDIX hydrolase	Tb927.11.15640 [23, 121, 122]
MERS2		MERS2	Targets MERS1 to RNA	PPR	Tb11.02.5120 [23]
Modification of the 3' end: Kinetoplast polyadenylation complex (KPAC)					
KPAP1		KPAP1	Major poly(A) polymerase	NT/TUTase, PAP associated	Tb927.11.7960 [31]
KPAF1	PPR1	KPAF1	mRNA A/U-tailing	PPR	Tb927.2.3180 [30, 170, 171]
KPAF2		KPAF2	mRNA A/U-tailing	PPR	Tb927.11.14380 [30]
KPAF3		KPAF3	mRNA stabilization/A-tailing	PPR	Tb927.9.12770 [29]
KPAF4		KPAF4	Poly(A) binding protein	PPR	Tb927.10.10160 [32]
U-insertion/deletion mRNA Editing: RNA editing catalytic complex (RECC)					
REN1	KREPB1	KREN1	U-deletion endonuclease	RNase III, PUF, ZF-C2H2	Tb927.1.1690 [86, 172]
REN2	KREPB3	KREN2	U-insertion endonuclease	RNase III, PUF, ZF-C2H2	Tb927.10.5440 [86, 173]
REN3	KREPB2	KREN3	U-insertion endonuclease	RNase III, PUF, ZF-C2H2	Tb927.10.5320 [85, 86, 98]
REX1	KREX1	KREX1	3'-5' U-specific exonuclease	Exo/endo/phos (EEP)	Tb927.7.1070 [85, 86, 97]
REX2	KREX2	KREX2	3'-5' U-specific exonuclease	Exo/endo/phos (EEP)	Tb927.10.3570 [85, 86, 97]
RET2	KRET2	KRET2	U-insertion TUTase	TUTase, PAP associated	Tb927.7.1550 [48, 85, 86, 113]
REL1	KREL1	KREL1	RNA ligase (U-deletion)	RNA lig/RNL2	Tb927.9.4360 [47, 85, 86, 90]
REL2	KREL2	KREL2	RNA ligase (U-insertion)	RNA lig/RNL2	Tb927.1.3030 [85, 86, 90]
MP81	KREPA1	KREPA1		ZF-C2H2, OB fold	Tb927.2.2470 [85, 86, 174]
MP63	KREPA2	KREPA2		ZF-C2H2, OB fold	Tb927.10.8210 [85, 86, 175]
MP42	KREPA3	KREPA3		ZF-C2H2, OB fold	Tb927.8.620 [85, 86, 92, 176]
MP24	KREPA4	KREPA4		OB fold	Tb927.10.5110 [85, 86, 94]
MP19	KREPA5	KREPA5		OB fold	Tb927.8.680 [86]
MP18	KREPA6	KREPA6		OB fold	Tb927.10.5120 [85, 86, 91, 177]
MP46	KREPB4	KREPB4		RNase III, PUF, ZF-C2H2	Tb927.11.2990 [85, 86, 103]
MP44	KREPB5	KREPB5		RNase III, PUF, ZF-C2H2	Tb927.11.940 [85, 86, 176, 178]
MP49	KREPB6	KREPB6		RNase III, ZF-C2H2	Tb927.3.3990 [85, 86, 95, 102]
MP47	KREPB7	KREPB7		RNase III, ZF-C2H2	Tb927.9.5630 [95, 102, 104]
MP41	KREPB8	KREPB8		RNase III, ZF-C2H2	Tb927.8.5690 [95, 102, 104]
	KREPB9	KREPB9		RNase III, ZF-C2H2	Tb927.9.4440 [179, 180]
	KREPB10	KREPB10		RNase III, ZF-C2H2	Tb927.8.5700 [179, 180]
MEAT1		MEAT1	RECC-like associated TUTase	TUTase, PAP associated	Tb927.1.1330 [181]

U-insertion/deletion mRNA Editing: RNA editing substrate binding complex (RESC)						
GRBC1	GAP2	RESC1	gRNA binding/stabilization		Tb927.7.2570	[121, 122]
GRBC2	GAP1	RESC2	gRNA binding/stabilization		Tb927.2.3800	[121, 122]
GRBC3	MRB8620	RESC3			Tb927.11.16860	[52, 124, 129]
GRBC4	MRB5390	RESC4			Tb11.02.5390b	[52, 129, 182]
GRBC5	MRB11870	RESC5			Tb927.10.11870	[52, 129, 183]
GRBC6	MRB3010	RESC6			Tb927.5.3010	[52, 123, 129, 133, 184]
GRBC7	MRB0880	RESC7			Tb927.11.9140	[52, 129]
REMC1	MRB10130	RESC8	RNA binding	ARM/HEAT repeats	Tb927.10.10130	[52, 120, 125, 129]
REMC2	MRB1860	RESC9			Tb927.2.1860	[52, 129]
REMC3	MRB800	RESC10			Tb927.7.800	[52, 129]
REMC4	MRB8180 MRB4150	RESC11A RESC11B	RNA binding		Tb927.8.8180 Tb927.4.4150	[52, 78, 129]
REMC5	MRB4160	RESC12	RNA binding		Tb927.4.4160	[52, 128, 129, 185]
REMC5 A	MRB8170	RESC12A			Tb927.8.8170	[52, 78, 128, 129, 185]
TbRGG2	TbRGG2	RESC13	RNA binding	RGG, RRM	Tb927.10.10830	[52, 78, 127, 129, 182, 186-188]
	MRB7260	RESC14		PhyH	Tb927.9.7260	[126, 129]
PAMC1		RESC15			Tb927.1.1730	[52]
PAMC2		RESC16			Tb927.6.1200	[52]
PAMC3		RESC17			Tb927.10.1730	[52]
PAMC4		RESC18			Tb927.1.3010	[52]
MERS3	RBP7910	RESC19		Z-DNA binding	Tb927.10.7910	[23, 131]
U-insertion/deletion mRNA Editing: REH2 RNA Helicase Complex (REH2C)						
REH2		KREH2	RNA helicase, RNA binding	DEAH/RHA, HA2, DSRM, OB/NTP_binding	Tb927.4.1500	[120-123, 132-135]
H2F1	MRB1680	KH2F1		ZF-C2H2	Tb927.6.1680	[134, 135, 182]
H2F2		KH2F2			Tb927.6.2140	[132, 134, 135]
Auxiliary RNA Processing Factors						
KPAP2		KPAP2	Putative poly(A) polymerase	NT/TUTase, PAP associated	Tb927.10.160	[136]
REH1	mHEL61	KREH1	RNA helicase	DEAD/DEAH box helicase	Tb927.11.8870	[139, 189]
MRP1	gBP21	KMRP1	RNA binding		Tb927.11.1710	[130, 140-148]
MRP2	gBP25	KMRP2	RNA binding		Tb927.11.13280	[130, 142, 143, 146-148]
RGG1		KRGG1	RNA binding		Tb927.6.2230	[122, 156, 157]
RBP16		KRBP16	RNA binding	Cold-shock RNA binding	Tb927.11.7900	[146, 148-155]
	MRB1590	KRBP72	RNA binding	ABC-like ATPase domain	Tb927.3.1590	[159]
TbRGG3	MRB1820	KRGG3			Tb927.3.1820	[129, 158]
REAP-1		KREAP1	RNA binding		Tb927.10.9720	[160-162]
RND		KRND1	U-specific 3'-5' exonuclease	RND, ZF-C2H2	Tb927.9.12720	[164]
PRORP2		KRNP1	RNase P	PRORP, PPR	Tb927.11.3010	[165]
mRPN1		KRPN1	Endonuclease	RNase III	Tb927.11.8400	[42, 46]

1089

1090

1091 **Figure 1.** A schematic diagram of mitochondrial RNA processing in *T. brucei*. The flow of
1092 processing reactions does not imply an experimentally established timing of these events. For
1093 example, the rRNA assembly into the ribosome or 5' pyrophosphate removal from mRNA
1094 may occur co-transcriptionally. Likewise, the mRNA may be edited prior to completion of 3'-
1095 5' trimming and 3' adenylation.




Fig. 1